




# Review Article: An Overview on Novel Antioxidant and Anti-Cancer Properties of Lycopene: A Comprehensive Review



Manjeet Gupta<sup>1</sup> , Mujeebullah Panizai<sup>2</sup>, Muhammad Farooq Tareen<sup>2</sup>, Sylvia Ortega-Martinez<sup>3</sup> , Nanuli Doreulee<sup>4\*</sup> 

1. Developmental Toxicology Laboratory, CSIR-Indian Institute of Toxicology Research, Lucknow, India.

2. Livestock & Dairy Development Department Balochistan, Quetta, Pakistan.

3. Department of Neurobiology, University of Chicago, Chicago, IL, USA.

4. Professor of Biological Sciences, Faculty of Exact and Natural Sciences, Head of direction of Human and Animal Physiology, Tbilisi State University, Tbilisi, Georgia.



**Citation** Gupta M, Panizai M, Farooq Tareen M, Ortega-Martinez S. An Overview on Novel Antioxidant and Anti-Cancer Properties of Lycopene: A Comprehensive Review. GMJ Medicine. 2018; 2(1):45-50. <https://doi.org/10.29088/GMJM.2018.45>

**doi** <https://doi.org/10.29088/GMJM.2018.45>



## Article info:

**Received:** 10 June 2018

**Accepted:** 17 August 2018

**Available Online:** 10 September 2018

**Checked for Plagiarism:** Yes

**Peer reviewers approved by:**

Dr. Melika Andrew

**Language Editor:**

Prof. Dr. Muhammad Azam Kakar

**Editor who approved publication:**

Prof. Dr. Nanuli Doreulee

## Keywords:

Antioxidant, Cancers,  
Lycopene

## ABSTRACT

Lycopene, a non-provitamin A carotenoid, is the reason for redness in tomatoes and some other vegetables. Lycopene has also been known as one fat-soluble red pigment that is produced by plants and some microorganisms. Nowadays, the effects of lycopene on healthiness have recently been significantly interested. Antioxidant properties of lycopene have been received attention as an anticancer. Different studies have investigated the effects of lycopene in relation to different cancers types. Lycopene significantly shows powerful anticancer activity against prostate cancer, even in progressed and aggressive condition. This review article aims to introduce the lycopene and possible mechanisms for the treatment of cancer.

## Introduction

**L**ycopene is well known for one fat-soluble red pigment that is synthesized in some plants and microorganisms [1]. Lycopene is extensively found in tomatoes and insignificant amount in other plants such as

papaya [2, 3]. Lycopene is one lipophilic acyclic isomer of  $\beta$ -carotene but it did not perform activity of vitamin A [4] and that could be attributed to absence of a  $\beta$ -ionic ring structure. It is widely found in human blood plasma with low-density lipoprotein fractions [5, 6]. As the Lycopene is originated from  $\beta$ -carotene and  $\alpha$ -tocopherol,

\* Corresponding Author:

**Nanuli Doreulee, PhD.**

**Address:** Head of Direction of Human and Animal Physiology, Tbilisi State University, Tbilisi, Georgia.

**E-mail:** [nanuli.doreuli@tsu.ge](mailto:nanuli.doreuli@tsu.ge)

it shows significantly higher antioxidant activity and has been extensively interested. [7, 8].

Skin lycopene is known to have sensitive rather than UV light stress. Lycopene present in natural plant sources is broadly found in trans form that have the high thermodynamic stability [9]. However, all the trans, 5-cis, 9-cis, 13-cis and 15-cis are known as the most common isomeric forms in lycopene [9]. The biological importance of the isomers in lycopene has been not still elucidated. It has been accepted that lycopene present in tomatoes, natural trans form, is slowly absorbed. Previous studies have reported that heat processing of tomatoes and tomato products causes to induce the isomerization of lycopene to the cis form which elevates its bioavailability [10]. Some evidences have been reported that isomerization reactions may occur in the cells. For example, increased concentration of cis isomers in serum and prostate tissue implicates the in vivo isomerization of lycopene [9].

Importance of lycopene for the treatment and prevention of cancer has been accepted. Elevation of oxidative stress plays a significant role in cancer risk. Lycopene is known to have powerful antioxidant properties compared to other known different carotenoids [11], that is as follows lycopene >  $\alpha$ -tocopherol >  $\alpha$ -carotene >  $\beta$ -cryptoxanthin > zeaxanthin =  $\beta$ -carotene > lutein [12]. Lycopene has ability to decrease oxidative damage through stimulation of enzyme activity of antioxidant [13]. Lycopene also inhibits the oxidative injuries of macromolecules [14]. Lycopene is involved in immune system and induces cellular apoptosis [15]. It has also been suggested to prevent reactive oxygen species formation and reduced phosphorylation of extracellular signal-regulated kinase that helps in prevention of cancer cell growth [16, 17]. This paper aims to investigate the anti-cancer effects of lycopene, especially mechanisms.

### Chemistry of lycopene

Lycopene is one non-cyclic carotenoid which is found in tomatoe. Lycopene, one hydrocarbon carotenoid

with general formula of C<sub>40</sub>H<sub>56</sub> contains one acyclic open-chain structure that comprises 13 double-bonds (Figure 1).

The double bonds present in lycopene are exposed to isomerization and various cis isomers (mainly 5, 9, 13, or 15) are observed in plants and blood plasma [18]. Physical variables including heat, light or some chemical reactions could cause isomerization from the trans-isomer into different mono-or poly-cis structures [11]. There are non-conjugated (2 bonds) and conjugated double bonds (11 cases) that are known to have antioxidant properties in lycopene [19]. Since the body cannot produce carotenoids, it is essential to have lycopene by diet.

### Antioxidant activity of lycopene

Oxidative stress is known as one of reasons for increased risk of cancer. Lycopene has been reported to have antioxidant properties as shown by in vitro experimental systems [11]. It is found that carotenoids reactivity not only relies on molecular and physical compounds but it also depends on their location or place of action into cells, ability to interact with other antioxidants, concentration and the partial pressure of oxygen [20-22]. Antioxidant activity could be attributed to polyene structures that are rich in electrons. Lycopene has been known to have potent oxygen scavenging reagent between carotenoids, and thus it modulates reactions activated by free radicals such as OH<sup>•</sup> or peroxy radicals [23].

Lycopene and other carotenoids have been suggested for their antioxidant properties in order to prevention of free radical reactions. Peroxyl radicals are produced in the organism when lipid peroxidation occurs and that could cause damages in lipophilic parts. The carotenoid oxidation products are epoxides placed in the  $\beta$ -ionone ring, as well as those are in the central double bond of the conjugated polyene chain. The most products of reaction are included ketones and aldehydes in the  $\beta$ -ionone ring. Prevention of these radical reactions by lycopene can prevent membranes from lipid peroxidation [23].




Figure 1. Molecular structure of lycopene

Some studies have shown that lycopene could upregulate the antioxidant Electrophile/antioxidant Response Element (EpRE/ARE) and the nuclear Factor E2-related factor 2 (Nrf2). It stimulates the formation of phase II detoxifying antioxidant enzymes that keep the cells safe from reactive species [24]. It is shown that Nrf2 nuclear transcription pathway could upregulate ARE system in HepG2 and MCF-7 cells [24] and is also involved in expression of ARE-regulated proteins such as Epoxide Hydrolase 1 (EPHX1), Superoxide Dismutase-1 (SOD-1), Catalase (CAT), and the metal binding protein Transferrin (TF), in prostate cells [25].

Enzymatic activity by lycopene induces Nrf2-mediated expression of phase II detoxifying/antioxidant enzymes [26]. A damage in DNA was usually induced by H<sub>2</sub>O<sub>2</sub>. Studies have shown that high levels of H<sub>2</sub>O<sub>2</sub> caused to induce the damage in DNA and each cell could protect itself against damage [27]. In a study, subjects consuming tomato for 2 weeks showed a strong inverse association between plasma lycopene content and lymphocyte DNA damage. Bašt et al. has shown that lycopene may increase the cellular antioxidant defense system through increasing non-enzymatic antioxidants [28].

Vitamins E and C from their radicals and also reduces δ-tocopheryl radical. Agarwal et al. have shown a significant decrease of serum lipid and LDL oxidation when subjects consumed tomato [29]. In addition, some studies have shown a role of ROS for the Ras superfamily of small GTPase, in redox regulation and ROSs have been

known to have significant downstream effects for Ras protein.

It is found that lycopene is capable to rescue Ras activation through reducing its pharnesylation and also by conducting its translocation from the membrane to cytoplasm in cells of patients with cancer [14] and also in patients with stimulated macrophages [17]. Changed in Ras activation were severely associated with prevention of the expression of 3-Hydroxy-3-Methylglutaryl Coenzyme A (HMG-CoA) reductase through the carotenoid and were associated with reduced ROS formation and activation of MAPK/NF-κB. qqq

**Anti-cancer activity of lycopene**

In this section, we discuss anti-cancer mechanisms of lycopene. Carotenoids have been known to have direct role in some redox-sensitive signalling pathways, changed in cancer [30]. It is also known to have ability to modify Ras activation through reducing pharnesylation and also inducing translocation in the membrane into the cytoplasm in cancer cells [31] and also stimulating macrophages [32]. It has also been known to have ability for suppression of MAPK phosphorylation and NF-κB activation in prostatic cancer cells [32]. Lycopene prevents AP-1 signalling in mammary cells [33].

It is believed that lycopene inhibits cancer through induction of apoptosis [34]. Zhang et al. [35] have reported that lycopene and its auto-oxidant products could induce apoptosis in HL-60 cells. Lycopene is involved in Bcl-2




Figure 2. Anti-cancer mechanisms of lycopene

Downloaded from gmjpress.de at 0:03 +0330 on Wednesday September 29th 2021 [ DOI: 10.29088/GMJM.2018.45 ]

that modulates the apoptosis that could be attributed to antioxidant properties of lycopene [34]. Lycopene also inhibits carcinogenesis by rescuing essential biomolecules such as lipids, LDL, proteins and DNA [35]. It has ability to quench singlet oxygen that could be credited to its conjugated double bonds [7].

A study has been shown that lycopene prevented growth of human endometrial, mammary and lung cancer cells, grown in *in vitro* culture [36]. It keeps cells against microcystinCR-induced mouse hepatocarcinoma through suppression of the phosphorylation of regulatory proteins and arrest of cells in the G0/G1 phase of the cell cycle [36]. Other studies have shown that lycopene decreases cellular proliferation created through IGF-1 in the different cancer cell lines [36].

Previous studies have shown that use of lycopene prevented cell cycle progression by modulation in G0/G1 stage by down-regulation of IGF-1R expression and the following decreased cell cycle regulatory proteins, such as cyclin D1, cyclin E and Cyclin-Dependent Kinases (CDK) 2 and 4 in breast and prostate cancer cell lines [37-39]. Lycopene-induced prevention of DNA formation has been seen in HL-60 promyelocytic leukaemia cells by an (3H) thymidine incorporation evaluation and caused cell cycle arrest in the G0/G1 phase [40].

Lycopene has been known to have ability for attenuating the phenotypic and functional maturation of murine bone marrow-dendritic cells that are present in Lipopolysaccharide (LPS)-induced DC maturation through down-regulating the expression of costimulatory molecules (CD80 and CD86) and significant histocompatibility complex type II molecules and through prevention of activation of MAPK and NF- $\kappa$ B [41]. Some papers have been reported the assess the effect of lycopene and/or tomato consumption, in the involvement as the markers of oxidative stress and on changed redox signalling. Some believed that lycopene shows anti-cancer properties by modulation in antioxidant properties.

Devaraj et al. have been examined the antioxidant ability of 8 weeks of lycopene consumption (6.5, 15, 30 mg/d) after 14 days washout period in healthy individuals [42]. However, their results showed that used doses could not have significant effect on LDL oxidation rate, plasma lipid peroxidation markers, and urinary F2-isoprostanes. They also showed that consumption of 30 mg/d dose could alleviate the lymphocyte DNA damage and urinary 8-OHdG contents in comparison to baseline levels. Another study has been shown that lycopene supplementation could not change total antioxidant

capacity or oxidized-LDL antibody levels, but it reduced serum MDA levels in comparison to baseline levels [43]. A summary of Anti-cancer of lycopene is shown in Figure 2.

## Conclusion

Consumption of lycopene has been relatively related with a decrease in cancer risk. The pharmacokinetic properties of lycopene for prevention and treatment of cancers have been cleared. We recommend the use of lycopene for treatment and prevention of cancer.

## Ethical Considerations

### Compliance with ethical guidelines

There was no ethical considerations to be noted in this article.

### Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### Authors' contributions

All authors contributed in data analysis, drafting and revising the paper and agreed to be responsible for all aspects of this work.

### Conflict of interest

The authors declared no conflict of interest.

## References

- [1] Krubasik P, Sandmann G. A carotenogenic gene cluster from *Brevibacterium linens* with novel lycopene cyclase genes involved in the synthesis of aromatic carotenoids. *Mol Gen Genet.* 2000; 263: 423-432. DOI: [DOI:10.1007/s004380051186](https://doi.org/10.1007/s004380051186)
- [2] Chandrika UG, Fernando KS, Ranaweera KK. Carotenoid content and *in vitro* bioaccessibility of lycopene from guava (*Psidium guajava*) and watermelon (*Citrullus lanatus*) by high-performance liquid chromatography diode array detection. *Int J Food Sci Nutr.* 2009; 60:558-566. [\[DOI:10.3109/09637480801987195\]](https://doi.org/10.3109/09637480801987195) [PMID]
- [3] Maiani G, Cañon MJ, Catasta G, Toti E, Cambrodon IG, Bysted A, et al. Carotenoids: actual knowledge on food sources, intakes, stability and bioavailability and their protective role in humans. *Mol Nutr Food Res.* 2009; 53(Suppl 2):194-218. [\[DOI:10.1002/mnfr.200800053\]](https://doi.org/10.1002/mnfr.200800053) [PMID]

- [4] Rao AV, Agarwal S. Role of antioxidant lycopene in cancer and heart disease. *J Am Coll Nutr*. 2000;19:563-569. [DOI:10.1080/07315724.2000.10718953] [PMID]
- [5] Clinton SK. Lycopene: chemistry, biology, and implications for human health and disease. *Nutr Rev*. 1998;56:35-51. [DOI:10.1111/j.1753-4887.1998.tb01691.x] [PMID]
- [6] Erdman JW, Jr, Bierer TL, Gugger ET. Absorption and transport of carotenoids. *Ann N Y Acad Sci*. 1993;691: 76-85. [DOI:10.1111/j.1749-6632.1993.tb26159.x] [PMID]
- [7] Di Mascio P, Kaiser S, Sies H. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch Biochem Biophys*. 1989;274:532 - 538. [DOI:10.1016/0003-9861(89)90467-0]
- [8] Miller NJ, Sampson J, Candeias LP, Bramley PM, Rice-Evans CA. Antioxidant activities of carotenes and xanthophylls. *FEBS Lett*. 1996;384:240-242. [DOI:10.1016/0014-5793(96)00323-7]
- [9] Clinton SK, Emenhiser C, Schwartz SJ, Boštwick DG, Williams AW, Moore BJ, Erdman JW: Cis-trans lycopene isomers, carotenoids and retinol in the human prostate. *Cancer Epidemiol Biomarkers Prev*. 1996; 5:823-833.
- [10] Stahl W, Sies H: Uptake of lycopene and its geometrical isomers is greater from heat-processed than from unprocessed tomato juice in humans. *J Nutr*. 1992; 122:2161-2166. [DOI:10.1093/jn/122.11.2161] [PMID]
- [11] Di Mascio P, Kaiser S, Sies H. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch Biochem Biophys*. 1989; 274:532-8. [DOI:10.1016/0003-9861(89)90467-0]
- [12] Heber D, Lu QY. Overview of mechanisms of action of lycopene. *Exp Biol Med*. 2002;227:920-3. [DOI:10.1177/153537020222701013] [PMID]
- [13] Velmurugan B, Bhuvaneshwari V, Burra UK, Nagini S. Prevention of N-methyl-N'-nitro-N-nitrosoguanidine and saturated sodium chloride-induced gastric carcinogenesis in Wistar rats by lycopene. *Eur J Cancer Preve*. 2002;11:19-26. [DOI:10.1097/00008469-200202000-00004]
- [14] Palozza P, Simone R, Catalano A, Boninsegna A, Böhm V, Fröhlich K, et al. Lycopene prevents 7-ketocholesterol-induced oxidative stress, cell cycle arrest and apoptosis in human macrophages. *J Nutr Biochem*. 2010;21:34-46. [DOI:10.1016/j.jnutbio.2008.10.002] [PMID]
- [15] Rao AV, Ray MR, Rao LG. Lycopene. *Adv Food Nutr Res*. 2006; 51:99-164. [DOI:10.1016/S1043-4526(06)51002-2]
- [16] Rao LG, Mackinnon ES, Josse RG, Murray TM, Strauss A, Rao AV. Lycopene consumption decreases oxidative stress and bone resorption markers in postmenopausal women. *Osteoporosis Int*. 2007;18:109-115. [DOI:10.1007/s00198-006-0205-z] [PMID]
- [17] Palozza P, Colangelo M, Simone R, Catalano A, Boninsegna A, Lanza P, et al. Lycopene induces cell growth inhibition by altering mevalonate pathway and Ras signaling in cancer cell lines. *Carcinogen*. 2010; 31:1813-1821. [DOI:10.1093/carcin/bgq157] [PMID]
- [18] Canene-Adams K, Campbell JK, Zaripheh S, Jeffery EH, Erdman JW. The tomato as a functional food. *J Nutr*. 2005; 135: 1226-1230. [DOI:10.1093/jn/135.5.1226] [PMID]
- [19] Shi J and Le Maguer M. Lycopene in tomatoes: Chemical and physical properties affected by food processing. *Crit Rev Biotechnol*. 2000; 20: 293-334. [DOI:10.1080/07388550091144212] [PMID]
- [20] Raiola A, Rigano MM, Calafiore R, Frusciante L, Barone A. Enhancing the Health-Promoting Effects of Tomato Fruit for Biofortified Food. *Med Inflamm*. 2014; 1-16. <http://dx.doi.org/10.1155/2014/139873>. [DOI:10.1155/2014/139873] [PMID] [PMCID]
- [21] Krinsky NI. Antioxidant functions of carotenoids. *Free Radic Biol Med*. 1989; 7 (6): 617-635. [DOI:10.1016/0891-5849(89)90143-3]
- [22] Frusciante L, Carli P, Ercolano MR. Antioxidant nutritional quality of tomato. *Moi Nutr Food Res*. 2007; 51: 609-617. [DOI:10.1002/mnfr.200600158] [PMID]
- [23] Stahl W, Sies H. Antioxidant activity of carotenoids. *Moi Asp Med*. 2003; 24: 345-351. [DOI:10.1016/S0098-2997(03)00030-X]
- [24] Ben-Dor A, Steiner M, Gheber L, Danilenko M, Dubi N, Linnewiel K, Zick A, Sharoni Y, Levy J. Carotenoids activate the antioxidant response element transcription system. *Mol Cancer Ther*. 2005; 4: 177-186.
- [25] Goo YA, Li Z, Pajkovic N, Shaffer S, Taylor G, Chen J, et al. Systematic investigation of lycopene effects in LNCaP cells by use of novel large-scale proteomic analysis software. *Proteomics Clin App*. 2007; 1: 513-523. [DOI:10.1002/prca.200600511] [PMID] [PMCID]
- [26] Linnewiel K, Ernst H, Caris-Veyrat C, Ben-Dor A, Kampf A, Salman H, et al. Structure activity relationship of carotenoid derivatives in activation of the electrophile/ antioxidant response element transcription system. *Free Radic Biol Med*. 2009; 47: 659-667. [DOI:10.1016/j.freeradbiomed.2009.06.008] [PMID]
- [27] Duthie SJ, Ma A, Ross MA, Collins AR. Antioxidant supplementation decreases oxidative DNA damage in human lymphocytes. *Cancer Res*. 1996; 56:1291-5.
- [28] Bašt A, Haenen GR, van den Berg R, van den Berg H. Antioxidant effects of carotenoids. *Int J Vitam Nutr Res*. 1998; 68:399-403. PMID: 9857268.
- [29] Agarwal A, Shen H, Agarwal S, Rao AV. Lycopene content of tomato products: its stability, bioavailability and in vivo antioxidant properties. *J Med Food*. 2001; 4:9-15. [DOI:10.1089/10966200152053668] [PMID]
- [30] Palozza P, Parrone N, Catalano A, Simone R. Tomato lycopene and inflammatory cascade: basic interactions and clinical implications. *Curr Med Chem*. 2010; 17: 2547-2563. [DOI:10.2174/092986710791556041] [PMID]
- [31] Palozza P, Colangelo M, Simone R, Catalano A, Boninsegna A, Lanza P, et al. Lycopene induces cell growth inhibition by altering mevalonate pathway and Ras signaling in cancer cell lines. *Carcinogenesis*. 2010; 31: 1813-1821. [DOI:10.1093/carcin/bgq157] [PMID]
- [32] Palozza P, Simone R, Catalano A, Boninsegna A, Böhm V, Fröhlich K, et al. Lycopene prevents 7-ketocholesterol-induced oxidative stress, cell cycle arrest and apoptosis in human macrophages. *J Nutr Biochem*. 2010; 21: 34-46. [DOI:10.1016/j.jnutbio.2008.10.002] [PMID]
- [33] Karas M, Amir H, Fishman D, Danilenko M, Segal S, Nahun A, et al. Lycopene interferes with cell cycle progression and insulin-like growth factor I signalling in mammary cancer cells. *Nutr Cancer*. 2000; 36: 101-111. [DOI:10.1207/S15327914NC3601\_14] [PMID]
- [34] Agarwal S, Rao AV. Tomato lycopene and low density lipoprotein oxidation: a human dietary intervention study. *Lipids*. 1998; 33:981-984. [DOI:10.1007/s11745-998-0295-6] [PMID]

- [35] Palozza P, Serini S, Di Nicuolo F, Calviello G. Modulation of Apoptotic Signalling by Carotenoids in Cancer Cells. *Arch Biochem Biophys.* 2004; 430: 104-109. [DOI:10.1016/j.abb.2004.02.038] [PMID]
- [36] Levy J, Bosin E, Feldmen B, Giat Y, Miinster A, Danilenko M, Sharoni Y. Lycopene is a more potent inhibitor of human cancer cell proliferation than either a-carotene or b-carotene. *Nutr Cancer.* 1995; 24:257-266. [DOI:10.1080/01635589509514415] [PMID]
- [37] Nahum A, Hirsch K, Danilenko M, Watts CK, Prall OW, Levy J, Sharoni Y. Lycopene inhibition of cell cycle progression in breast and endometrial cancer cells is associated with reduction in cyclin D levels and retention of p27(Kip1) in the cyclin E-cdk2 complexes. *Oncogene.* 2001;20: 3428-3436. [DOI:10.1038/sj.onc.1204452] [PMID]
- [38] Nahum A, Zeller L, Danilenko M, Prall OW, Watts CK, Sutherland RL, et al. Lycopene inhibition of IGF-induced cancer cell growth depends on the level of cyclin D1. *Eur J Nutr.* 2006; 45:275-282. [DOI:10.1007/s00394-006-0595-x] [PMID]
- [39] Siler U, Barella L, Spitzer V, Schnorr J, Lein M, Goralczyk R, Wertz K. Lycopene and vitamin E interfere with autocrine/ paracrine loops in the Dunning prostate cancer model. *FASEB J.* 2004;18:1019-1021. [DOI:10.1096/fj.03-1116fje] [PMID]
- [40] Ivanov NI, Cowell SP, Brown P, Rennie PS, Guns ES, Cox ME. Lycopene differentially induces quiescence and apoptosis in androgen-responsive and -independent prostate cancer cell lines. *Clin Nutr.* 2007; 26:252-263. [DOI:10.1016/j.clnu.2007.01.002] [PMID]
- [41] Kim GY, Kim JH, Ahn SC, Lee HJ, Moon DO, Lee CM, Park YM. Lycopene suppresses the lipopolysaccharide-induced phenotypic and functional maturation of murine dendritic cells through inhibition of mitogen-activated protein kinases and nuclear factor- $\kappa$ B. *Immunol.* 2004; 113: 203-211. [DOI:10.1111/j.1365-2567.2004.01945.x] [PMID] [PMCID]
- [42] Devaraj S, Mathur S, Basu A, Aung HH, Vasu VT, Meyers S, Jialal I. A dose-response study on the effects of purified lycopene supplementation on biomarkers of oxidative stress. *J Am Coll Nutr.* 2008; 27: 267-273. [DOI:10.1080/07315724.2008.10719699] [PMID] [PMCID]
- [43] Neyeštani TR, Shariatzadeh N, Gharavi A, Kalayi A, Khalaji N. Physiological dose of lycopene suppressed oxidative stress and enhanced serum levels of immunoglobulin M in patients with Type 2 diabetes mellitus: a possible role in the prevention of long-term complications. *J Endocrinol Invest.* 2007; 30: 833-838. [DOI:10.1007/BF03349224] [PMID]