




The Effects of Oral Administration of Pulegone in Carbon Tetrachloride-Induced Oxidative Stress in Wistar Rats

Antonina Rabinovich¹, Nina Romanoff^{1*}, Dominika Mordvinov², Margarita Ivanov²



¹Pirogov Russian National Research Medical University, Moscow, Russia;
²Institute for Biomedical Research, Association of Pharmaceutical Research and Development, Kyiv, Ukraine.

•Corresponding Author.

Nina Romanoff, Ph.D, Pirogov Russian National Research Medical University, Moscow, Russia.

✉: nromanoff@gmail.com

Received. 15 June, 2019

Accepted. 10 August, 2019

Published. 25 August, 2019

Checked for Plagiarism. Yes

Peer reviewers approved by:

Dr. Melika Andrew

Language Editor:

Prof. Dr. Mohammad Azam Kakar

Editor who approved publication:

Prof. Dr. Nanuli Doreulee

Abstract

Background and purpose. Carbon tetrachloride (CCl₄) has been applied to induce the toxicity and hepatic fibrosis. Natural antioxidants are known as efficient and safe treatments for hepatotoxicity compared with synthetic antioxidants. This study aimed to evaluate the effects of oral administration of pulegone in carbon tetrachloride-induced oxidative stress in Wistar rats.

Materials and Methods. Twenty rats were randomly assigned into four groups including control animals that received olive oil, Toxic control that administrated with 30% CCl₄, Pulegone-20 & 30 that administrated with pulegone 20 mg/kg and 30 mg/kg, respectively, with in along to 30% CCl₄. The liver levels of total cholesterol, triacylglycerides, phospholipids, superoxide dismutase (SOD), catalase (CAT), reduced glutathione (GSH), glutathione peroxidase (GPx), and vitamins C and E were evaluated.

Results. Administration of CCl₄ increased levels of cholesterol, triacylglycerides and lipid oxidation, but it also reduced levels of phospholipids, SOD, CAT, GSH, GPx, and vitamins C and E (P<0.05). Oral administration of pulegone, especially in the higher levels, could reverse negative effects of CCl₄ (P<0.05).

Conclusion. Using pulegone is recommended for liver protection due to its vital therapeutic antioxidant properties.

Keywords. Animal model, Enzymatic antioxidants, Liver cholesterol, Vitamin C



Introduction

Carbon tetrachloride, CCl₄ is usually applied to induce the toxicity and hepatic fibrosis. The hepatotoxic effect of CCl₄ could be attributed to its fast cleavage through cytochrome P450 in hepatocytes, that produces trichloromethyl radicals leading to lipid peroxidation and then to membrane injury [1]. Activation of the cytochrome catabolizes CCl₄ to trichloromethyl radical (CCl₃·) which produces toxic reactive trichloromethyl peroxy radical [2]. Oxidative stress increases in case of imbalance between formation and scavenging of reactive oxygen species (ROS) [2]. Oxidative stress is known as cause for the different degenerative diseases such as oxidative stress [3]. Natural antioxidants are known as efficient and safe treatments for hepatotoxicity compared with synthetic antioxidants [4, 5].

In this connection, Pulegone is a natural monoterpene ketone which is contained in the essential oil of a class of plants including *Mentha* species. It is reported that *Mentha pulegium* L. and *Mentha longifolia* L. have 60%–90% and 17% pulegone, respectively, as main component and also have anti-bacterial, antioxidant and anti-inflammatory properties [6, 7]. It is known to have antimicrobial activity, especially against all the *Salmonella* species [8]. A study showed that major volatile compounds of *Satureja macrostema* such as pulegone showed antioxidant activity in 2, 2-diphenyl-1-picrylhydrazyl and 2, 2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid [9]. It seems that pulegone could alleviate adverse effects of oxidative stress in carbon tetrachloride-induced oxidative stress. Thus, this study, for first time, was conducted to evaluate the effects of oral administration of pulegone in carbon tetrachloride-induced oxidative stress in Wistar rats.

Materials and methods

Animals

A total of 20 adult Male Wistar rats with initial weight of 220±20 g, were used in this study. Animals had allowed to ad libitum to a pelleted diet and water in 25±2 °C with a 12-h light/dark cycle. We did not use female rats to avoid the data variability obtained by hormonal cycles in females.

Experimental procedure

The Pulegone was procured from Sigma Chemical Co, USA. Based on previous acute toxicological reports of pulegone, 20 mg/kg body weight of pulegone was prepared to be nontoxic for rats [10]. The levels of 20 and 30 mg/kg doses of pulegone were used. Twenty rats were randomly allocated into four groups and five animals in per group and pulegone was gavaged with animal oral feeding cannula and divided as follows,

Group 1: control- olive oil (1 ml/kg), (Control),
Group 2: toxic control-30% CCl₄, (Toxic), Group 3:
Pulegone 20 mg/kg + 30% CCl₄, (Pulegone-20),
Group 4: Pulegone30 mg/kg + 30% CCl₄, (Pulegone-30)

Rats in Groups 2, 3, and 4 were treated with 30% CCl₄ olive oil (1 mL/Kg) by using intra-peritoneal administration for one time every three days for eight-week period [2]. Animals in pulegone groups received 20 mg/kg and 30 mg/kg pulegone, 72 h prior to CCl₄ treatment. In the end of trial, experimental animals were anesthetized and sacrificed by cervical dislocation, and the liver samples were collected, rinsed with saline solution, and stored at -80°C.

Investigation of lipid peroxidation, lipid profile, enzymatic and non-enzymatic antioxidants

Malondialdehyde (MDA) was assessed as one standard procedure for lipid peroxidation in the 532 nm as reported by others [11]. We have used Chloroform-methanol mixture (2:1 v/v) [12] for

extraction of lipids in the liver tissue for investigation of the total cholesterol [13], triacylglycerides [14, 15], and phospholipids [16]. Standard methods were applied to estimation of the superoxide dismutase (SOD) [17], catalase (CAT) [18], reduced glutathione (GSH), glutathione peroxidase (GPx) [19], Vitamin C [20] and vitamin E [21].

Statistical analysis

The data were analyzed by one-way Analysis of Variance (ANOVA) and Duncan's Multiple Range Test (DMRT). Statistical Package for Social Sciences (SPSS) 23.0 (IBM Corporation, Chicago, IL, USA) was used for analysis of the data and Graph Pad Prism was used for analysis of the data. The data were reported as means \pm SD.

Results

Effects of the different levels of pulegone on lipid peroxidation, Triacylglycerides, cholesterol and phospholipids in liver of Wistar rats are shown in Table 1. Results showed that levels of lipid peroxidation, triacylglycerides and cholesterol were significantly higher and also level of phospholipids was significantly lower in Toxic group compared with control group ($P < 0.05$). Oral administration of pulegone reversed adverse effects of CCl_4 on the mentioned parameters ($P < 0.05$). The best response was observed in level of 30 mg/kg pulegone. The data for enzymatic and non-enzymatic antioxidants are illustrated in Figure 1. As results show, CCl_4 lowered levels of antioxidants, but pulegone could partly spare antioxidants and the best response was observed in the highest level ($P < 0.05$).




Figure 1 Effects of the different levels of pulegone on enzymatic and non-enzymatic antioxidants in liver of rats. Superscripts (a-d) show significant differences among groups.

Table 1. Effects of the different levels of pulegone on lipid peroxidation (nmoles of MDA/mg protein), Triacylglycerides (mg/g tissue wt), cholesterol (mg/g tissue wt) and phospholipids (mg/g tissue wt) in liver of Wistar rats.

Parameters	Control	Toxic	Pulegone-20	Pulegone-30	P-values
Lipid peroxidation	1.19±0.05d	3.23±0.07a	2.83±0.15b	2.13±0.08c	**
Triacylglycerides	3.61±0.25d	6.59±0.17a	5.41±0.21b	4.83±0.13c	**
Cholesterol	3.21±0.08d	4.89±0.03a	4.30±0.05b	3.85±0.04c	**
Phospholipids	14.39±0.95a	11.21±0.77d	12.93±0.55c	13.21±0.38b	**

Superscripts (a-d) show significant differences among groups.

Discussion

The liver is known to have essential roles in metabolic homeostasis and it is also responsible for the metabolism, synthesis, storage and redistribution of nutrients and macromolecules. It is not only involved in the metabolism and detoxification of the body but also removes wastes and xenobiotics via metabolic conversion and biliary excretion [22]. The CCl₄ metabolism has been used as model for liver necrosis and fibrosis. Oxidative stress is known to have essential role in CCl₄-induced toxicity. Moreover, the induction of trichloromethyl-free radical increases lipid peroxidation process. It destroys membrane integrity and Ca²⁺ homeostasis for production of hepatocellular damage [23]. In the current study, the CCl₄ disturbed lipid profile and increased lipid oxidation and decreased levels of antioxidants. However, the use of pulegone could attenuate the effects of CCl₄ which could be attributed to its antioxidant activity. It is shown that antioxidant phyto-chemicals protect oxidative damages [24]. Natural antioxidants are involved in oxidative stress by ROS scavenging activity and induction of antioxidant and phase II detoxifying enzymes [25]. Improvement in lipids levels in pulegone could be attributed to CCl₄ mechanism. The CCl₄-induced liver injury stimulates lipid oxidation and increases production of ROS [23]. Our findings for lipid

oxidation showed higher levels in Toxic group in comparison to control group in terms of MDA, as final-product of membrane lipid oxidation. It means that increased lipid oxidation is paralleled decreased antioxidant enzymes, as illustrated in Figure 1. It seems that pulegone spares antioxidants and help to protect of the antioxidants. On the other hand, we observed a raised level in cholesterol and triacylglycerides in Toxic group. It is accepted that CCl₄ increases levels of cholesterol to rise in hepatocytes. On the other hand, reduced levels of phospholipids were seen following administration of CCl₄ treatment, that could be a result to an augmentation in phospholipase activity [26, 27]. To better understand of the pulegone activities, it is need to explain further the antioxidant parameters. GSH is one key non-enzymatic antioxidant which controls the intracellular redox homeostasis and protects cells from deleterious effects of ROS [28]. CCl₄ reduced levels of GSH but pulegone spared its levels. To be higher GSH levels can protect the liver against oxidative damage via directly scavenging the ROS by component of the GSH redox system including GPx, glutathione reductase, and glutathione-s-transferase [29]. On the other hand, results showed that CCl₄ reduced levels of vitamins C and E. Increased levels of vitamins means antioxidant nature of pulegone. It seems that

treatment with pulegone maintains membrane from damages and also protects active forms of vitamin C and vitamin E from ROS by increased levels of GSH, due to their close relation [30]. Three main enzymes CAT, GPx, and SOD are initial lines against damages. SOD is known to have ability for conversion of the superoxide anion to H₂O₂ and O₂. The H₂O₂ decreased is catalyzed via CAT and GPx that protects the tissue from biomolecule-damaging ROS [31]. Thus, CCl₄ reduces the levels of enzymatic and non-enzymatic antioxidants but pulegone attenuates its effects.

Conclusion

Taken together, the CCl₄ has adverse effects on liver lipids and increases levels lipid oxidation and also lowers the levels of antioxidants. Administration of pulegone alleviated the effects of CCl₄ on liver lipids and antioxidants. Pulegone spares the antioxidants and helps to protect against damages and thus reduces levels of lipids. It is recommended to use pulegone for treatment as it's a better choice for protection against liver damages.

Ethical Considerations

Compliance with ethical guidelines

Approval for this study was obtained from Association of Pharmaceutical Research and Development in Kyiv.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

All authors contributed toward data analysis, drafting and revising the paper and agreed to be responsible for all the aspects of this work.

Conflict of interest

The authors declared no conflict of interest.

References

1. El-Boshy ME, Abdelhamidb F, Richab E, AshshiaA, Gaiitha M, Qustya N. Attenuation of CCl₄ induced oxidative stress, immunosuppressive, hepatorenal damage by Fucoidan in rats. *J Clin Toxicol*. 2017; 7(3):1–7. [\[crossref\]](#)
2. Goodla L, h Manubolu M, Pathakoti K, Jayakumar T, Sheu J, Fraker M, Tchounwou PB, Poondamalli PR. Protective effects of ammannia baccifera against CCl₄induced oxidative stress in rats. *Int J Environ Res Public Health*. 2019; 16(8): 1440. [\[crossref\]](#)
3. Hensley K, Robinson KA, Gabbita SP, Salsman S, Floyd RA. Reactive oxygen species, cell signaling, and cell injury. *Free Radic Biol Med*. 2000; 28(10): 1456–1462. [\[crossref\]](#)
4. Kadri A, Zarai Z, Ben Chobba I, Bekir A, Gharsallah N, DamakM, Gdoura R. Chemical composition and antioxidant activity of Marrubium vulgare L. essential oil from Tunisia. *Afr J Biotechnol*. 2011; 10(19): 3908–3914.
5. Lavanya, G, Voravuthikunchai, S.P, Towatana, N.H. Acetone Extract from Rhodomyrtus tomentosa. A Potent Natural Antioxidant. *Evid Based Complementary Altern Med*. 2012; 2012: 535479. [\[crossref\]](#)
6. Jalilzadeh-Amin G, Maham M, Dalir-Naghadeh B, Kheiri F. Effects of Mentha longifolia essential oil on ruminal and abomasal longitudinal smooth muscle in sheep. *J Essent Oil Res*. 2012; 24: 61–69. [\[crossref\]](#)
7. Turner GW, Croteau R. Organization of monoterpene biosynthesis in Mentha. Immunocytochemical localizations of geranyl diphosphate synthase, limonene-6-hydroxylase, isopiperitenol dehydrogenase, and pulegone reductase. *Plant Physiol*. 2004; 136(4): 4215–4227. [\[crossref\]](#)

8. Flamini G, Cioni PL, Puleio R, Morelli I, Panizzi L. Antimicrobial activity of the essential oil of *Calamintha nepeta* and its constituent pulegone against bacteria and fungi. *Phytother Res*. 1999; 13(4): 349–351. [crossref]
9. Torres-Martínez R, García-Rodríguez YM, Ríos-Chávez P, Saavedra-Molina A, López-Meza JE, Ochoa-Zarzosa A, et al. Antioxidant activity of the essential oil and its major terpenes of *Satureja macrostema* (Moc. and Sessé ex Benth.) Briq. *Phycog Mag*. 2017; 13(4): S875–80. [crossref]
10. Cheraghali Z, Mohammadi R, Jalilzadeh-amin G. Planimetric and biomechanical study of local effect of pulegone on full thickness wound healing in rat. *Malays J Med Sci*. 2017; 24(5): 52–61. [crossref]
11. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem*. 1979; 95(2): 351–358. [crossref]
12. Folch J, Lees M, Stanley GHS. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem*. 1957; 226(1): 497–509. [PMID]
13. Parekh AC, Jung DH. Cholesterol determination with ferric acetate-uranium acetate and sulfuric acid-ferrous sulfate reagents. *Anal Chem*. 1970; 42(12): 1423–1427. [crossref]
14. Rice EW. Triglycerides in serum. In *Standards Methods in Clinical Chemistry*, Roedrick P, McDonal RP, Eds, Academic Press, New York, NY, USA, 1970, p. 215.
15. Van Handel E. Suggested modifications of the micro determination of triglycerides. *Clin Chem*. 1961; 7: 249–251. [PMID]
16. Rouser G, Fksischer S, Yamamoto A. Two dimensional thin layer chromatographic separation of polar lipids and determination of phospholipids by phosphorus analysis of spots. *Lipids*. 1970; 5(5): 494–496. [crossref]
17. Misra HP, Fridovich I. The Role of Superoxide Anion in the Autoxidation of Epinephrine and a Simple Assay for Superoxide Dismutase. *J Biol Chem*. 1972; 247(10): 3170–3175. [PMID]
18. Aebi H. Catalase in vitro. *Methods Enzymol*. 1984; 105: 121–126. [crossref]
19. Wendel A, Feuerstein S, Konz K-H. Acute paracetamol intoxication of starved mice leads to lipid peroxidation in vivo. *Biochem Pharmacol*. 1979; 28(13): 2051–2055. [crossref]
20. Omaye ST, Turnbull JD, Sauberlich HE. Selected methods for the determination of ascorbic acid in animal cells, tissues, and fluids. *Methods Enzymol*. 1979; 62: 3–11. [crossref]
21. Naziroglu M, Karaoglu A, Aksoy AO. Selenium and high dose vitamin E administration protects cisplatin-induced oxidative damage to renal, liver and lens tissues in rats. *Toxicol*. 2004; 195(2-3): 221–230. [crossref]
22. Joan O, Barbara AF, Qing X, Samuel WF. The identification of stem cells in human liver diseases and hepatocellular carcinoma. *Exp Mol Pathol*. 2010; 88(3): 331–340. [crossref]
23. Basu S. Carbon tetrachloride-induced lipid peroxidation: Eicosanoid formation and their regulation by antioxidant nutrients. *Toxicol*. 2003; 189(1-2): 113–127. [crossref]
24. Pathakoti, K, Goodla, L, Manubolu, M, Tencomnao, T. metabolic alterations and the protective effect of punicalagin against glutamate-induced oxidative toxicity in HT22 cells. *Neurotox Res*. 2017; 31(4): 521–531. [crossref]
25. Koolen HHF, da Silva FMA, Gozzo FbC, de Souza AQL, de Souza ADL. Antioxidant, antimicrobial activities and characterization of phenolic compounds from buriti (*Mauritia flexuosa* L. f.) by UPLC-ESI-MS/MS. *Food Res Int*. 2013; 51(2): 467–473. [crossref]
26. Lamb RG, Snyder JW, Coleman JB. New trends in the prevention of hepatocellular death. Modifiers of calcium movement and of membrane phospholipid metabolism. CRC Press: Boca Raton, FL, USA, 1988.

27. Coleman JB, Condie LW, Lamb RG. The influence of CCl₄ biotransformation on the activation of rat liver phospholipase C in vitro. *Toxicol Appl Pharmacol.* 1988; 95(2): 200–207. [[crossref](#)]
28. Miesel R, Sanocka D, Kurpisz M, Kroger H. Antiinflammatory effects of NADPH oxidase inhibitors. *Inflamm.* 1995; 19(3): 347–362. [[crossref](#)]
29. Aquilano K, Baldelli S, Ciriolo MR. Glutathione: New roles in redox signaling for an old antioxidant. *Front Pharmacol.* 2014; 5: 196. [[crossref](#)]
30. Winkler, B.S. Unequivocal evidence in support of the nonenzymatic redox coupling between glutathione/glutathione disulfide and ascorbic acid/dehydroascorbic acid. *Biochim Biophys Acta.* 1992; 1117(3): 287–290. [[crossref](#)]
31. Halliwell B, Gutteridge JM. *Free Radicals in Biology and Medicine*; Oxford University Press: Oxford, MI,USA, 2007.

GMJ Medical Press, LLC

Copyright. © 2019 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation. Rabinovich A, Romanoff N, Mordvinov D, Ivanov M. The Effects of Oral Administration of Pulegone in Carbon Tetrachloride-Induced Oxidative Stress in Wistar Rats. *GMJ Medicine.* 2019; 3: 132–138.

<https://doi.org/10.29088/GMJM.2019.132>