




Beneficial Effects of Curcumin on Rats with Polycystic Ovary Syndrome: Evaluation of the Gene Expression of GLUT4, $Er\alpha$ and Insulin Resistance

Mohamad Kazem Shahmoradi¹, Majid Askaripour², Soodeh Rajabi², Giga Dzigandzli^{3*}



¹Department of General Surgery, Lorestan University of Medical Sciences, Khorramabad, Iran;
²Department of Physiology, School of Medicine, Bam University of Medical Sciences, Bam, Iran;
³Georgian Center for Neuroscience Research, International Center for Intelligent Research, Tbilisi, Georgia.

*Corresponding Author, Giga Dzigandzli, MD, Georgian Center for Neuroscience Research, International Center for Intelligent Research, Tbilisi, Georgia.

✉ .gigadzizandzli@gmail.com

Received: 12 September, 2018

Accepted: 23 November, 2018

Published: 30 November, 2018

Checked for Plagiarism: Yes

Peer reviewers approved by:

Dr. Melika Andrew

Language Editor:

Prof Dr. Muhammad Azam Kakar

Editor who approved publication:

Prof. Dr. Nanuli Doreulee

Abstract

Background and purpose. Polycystic ovary syndrome (PCOS) is one of the most common endocrine disorders in the women and animals. Some common hormonal therapies are used to treat the PCOS condition. This study was conducted to evaluate the effects of curcumin on rats with PCOS by evaluation of the gene expression of the GLUT4 and $Er\alpha$ and insulin resistance.

Methods. Following induction of PCOS, sixty Sprague–Dawley female rats were divided into four groups including: (1) Control group (2) Control PCOS (3 & 4) those treated with 100 and 200 mg/kg of curcumin respectively. Body weight, fasting blood glucose (FBG), fasting insulin serum (FIS), homeostasis model assessment of insulin resistance (HOMA–IR) and gene expression of GLUT4 and $Er\alpha$ were evaluated.

Results. Induction of PCOS increased body weight, FBG, FIS, HOMA–IR and decreased gene expression of GLUT4 and $Er\alpha$ ($P < 0.05$), but oral administration of curcumin could alleviate adverse effects of PCOS on the mentioned parameters ($P < 0.05$).

Conclusion. It can be concluded that curcumin alleviates adverse effects of PCOS. It can be recommended to use the curcumin for the treatment of patients with PCOS.

Keywords. Curcumin, PCOS, $Er\alpha$, Fasting insulin, GLUT4, Rat

Introduction

Polycystic ovary syndrome (PCOS) is one of the most common endocrine disorders that induces anovulation in the women and animals [1]. Some metabolic abnormalities including dyslipidemia, insulin resistance, diabetes, obesity, and infertility occur in patients



with PCOS [2, 3]. PCOS has been known with some signs including production of small arrested antral follicle, decreased estrogen and increased LH/FSH ratio [4]. Previous studies have shown that insulin signaling pathway could have major role in pathogenesis of PCOS which comprises pathways of phosphatidylinositol 3-kinase (PI3K) and protein kinase B/Akt signaling [5-7]. Several studies have shown that selective insulin resistance can progress. In such cases, metabolic response to insulin action has been faulted including PI3K pathway, however, other actions are not only maintained but may be increased, including the mitogen-activated protein kinase (MAPK) pathway, in insulin group target tissues and ovary [5-6]. Insulin joins to its receptor and acts insulin receptor substrate (IRS) through the phosphorylation of tyrosine residues. The expression of IRS in the uterus has been reported [7] and glucose metabolism is essential for uterine cell differentiation [8]. It has been known that GLUT4 is the most important glucose transporter isoform in insulin-dependent tissues and modulates insulin-stimulated glucose transport in the uterus [9]. Different studies have reported decreased GLUT4 expression in the uterus of PCOS patients [10, 11]. Oestrogen receptor α (ER α) has been known to have an essential role [12], for example ER-knockout rats are infertile [13-15]. Some common hormonal therapies are used to treat the PCOS condition and to stimulate the ovulation. These treatments cause adverse effects on arthritis, joint or muscle pain [16] and psychological disturbances [17]. There is a great attention on medicines from natural sources that are safe and useful. Curcumin, is a natural active in turmeric, has biological effects including anti-inflammatory, antioxidant [18], hypoglycaemic [19] and anti-hyperlipidemic activities. It is known to have anti-proliferative and apoptotic activities in some human cancer cell lines, such as cells obtained from cancers of prostate, breast and ovary [20, 21]. Different published studies have showed its protective effects on porcine ovarian granulosa cells [22]. It seems that curcumin could improve the gene

expression of GLUT4 and ER α and other related parameters in rats with PCOS. Therefore, this study was conducted to evaluate the effects of curcumin on insulin resistance (IR) and uterine gene expression of GLUT4 and ER α in rats with PCOS.

Materials and methods

Animals

Sixty Sprague-Dawley female rats with weight of 180 ± 5 g were used for this study. All the animals were grouped in a 25°C a lighting diet 12:12 h light:dark cycle. The animals had ad libitum access to conventional feed pellets and water. To evaluate the body weight changes, we have weighed animals in start and end of the trial. All the animals were randomly grouped into four groups (n=15) including: 1) Control group without PCOS, 2) PCOS group without treatment, 3 & 4) PCOS rats treated with daily oral doses of 100 and 200 mg/kg, daily (Cur-100 and Cur-200). Curcumin was prepared from Sigma Chemicals Co, St. Louis, MO, USA. Curcumin was dissolved in 0.5% carboxy methyl cellulose per oral for 30 days. To induce the PCOS, animals were subcutaneously administrated with 6 mg dehydroepiandrosterone (DHEA) per 100 g day⁻¹ (DHEA dissolved in 0.2 mL sesame oil) for 21 consecutive days [23]. Following administration of curcumin (after 30 days), 5 animals per group were killed and some tissues including ovaries and uterus were collected and blood samples were collected to prepare the serum.

Evaluation of IR

The blood samples were used to evaluate the fasting blood glucose (FBG) and fasting insulin serum (FIS) levels. FBG was assessed by glucose oxidase procedure and FIS was assessed by a direct competitive enzyme-linked immunosorbent assay (ELISA) kit [24]. The optical values were read in the 450 nm by a microplate reader and the induction of IR was investigated by the homeostasis model assessment of insulin resistance (HOMA-IR) method.

HOMA-IR was calculated using the following formula:

$$\text{HOMA-IR} = \text{FBG (mmol/L)} \times \text{FIS (mU/L)} / 22.50.$$

Rat ovarian morphology

All the animals' ovaries were prepared by surgery fixed in 10% formalin and included in paraffin. Sections prepared were stained by haematoxylin and eosin then evaluated by two pathologists that did not information from the sample type for ovarian morphological properties.

Gene expression of GLUT4 and Era

Endometrium RNA isolation and real-time polymerase chain reaction (PCR) were conducted as reported by others [25]. The primers sequences were GLUT4, forward (5'- GGGCTGTGAGTGAGTGCTTTC-3') and reverse (5'- CAGCGAGGCAAGGCTAGA-3'); Era, forward (5'-CCAAAGCCTCGGGAATGG-3') and reverse (5'- AGCTGCGGGCGATTGAG-3'); and β -actin, forward (5'- AAGGCCAACCCTGAAAAGAT-3') and reverse (5'-ACCAGAGGCATACAGGGAC-3').

Statistical analysis

SPSS software was used to analysis the data. All the values are reported as mean \pm standard deviation (SD). The tukey test was used to investigate the difference between groups. The level of statistical significance was set at $P < 0.05$.

Results

Body weight

Effects of experimental treatments on body weight of rats with PCOS are shown in Figure 1. As results show, body weight was significantly higher in rats with PCOS in comparison to those in control group ($P < 0.05$). Oral administration of curcumin in the both levels (100 and 200 mg) significantly decreased adverse effects of PCOS on body weight

and better response was observed in higher dose ($P < 0.05$).






Figure 1 Effects of oral administration of curcumin on body weight (g) of rats with PCOS. Superscripts (a-d) show significant difference between groups.

Insulin resistance

Effects of oral administration of curcumin on FIS, FBG and HOMA-IR are shown in Figure 2. As results show, PCOS increased levels of FIS, FBG and HOMA-IR in comparison to control group ($P < 0.05$). Oral administration of curcumin could significantly decrease levels of FIS, FBG and HOMA-IR ($P < 0.05$).






Figure 2 Effects of oral administration of curcumin on insulin resistance of rats with PCOS. Superscripts (a-d) show significant difference between groups.

Ovarian morphology

Pathological evaluation and light microphotography of ovarian morphology are shown in Figure 3. In the control group, the follicles were normal, but ovaries of rats in the PCOS group had an increase in ovarian volume, decreased corpus luteum and theca layer hypertrophy and thickening in comparison to control group. Cystic follicles were also observed in PCOS group that means induction of PCOS. Oral administration of curcumin alleviated cystic follicles and increased corpus luteum.




Figure 3 Effects of oral administration of curcumin on ovarian morphology of rats with PCOS.

Gene expression

Effects of oral administration of curcumin on gene expression of GLUT4 and $Er\alpha$ of rats with PCOS are shown in Figure 4. Induction of PCOS decreased expression of GLUT4 and $Er\alpha$ ($P < 0.05$). The use of higher levels of curcumin increased expression of GLUT4 and $Er\alpha$ ($P < 0.05$). The use of lower level could also increase GLUT4 and $Er\alpha$ expression ($P < 0.05$).




Figure 4 Effects of oral administration of curcumin on gene expression of GLUT4 and $Er\alpha$ of rats with PCOS. Superscripts (a-d) show significant difference between groups.

Discussion

PCOS is one of the disorder related with diabetes and causes with signs such as hyperglycemia in first stages that gradually induces IR [26]. In the current study, rats induced with PCOS indicated an increase in FIS, FBG and HOMA-IR. Hyperglycaemia is result from IR, but the exact mechanism of PCOS is still not known. Previous studies have shown faulted insulin intracellular signaling, particularly alterations in insulin receptor substance-1 (IRS-1) phosphorylation that may be a factor for PCOS-IR [27]. On the other hand, adipose tissue increases the release of pro-inflammatory cytokines [28] and changes IRS-1 tyrosine phosphorylation to serine phosphorylation, which initiates IR. Our findings also showed that the body weight was significantly higher in PCOS rats. It seems overweight promotes to produce pro-inflammatory cytokines and promotes IR. Oral administration of curcumin improved IR, especially in higher doses. Improved IR could be attributed to anti-inflammatory properties

of curcumin that prevents production of pro-inflammatory cytokines [18]. On the other hand, curcumin decreased body weight that in turn decreases IR. As results showed, oral administration of curcumin decreased levels of FBG and FIS. Decreased levels of FBG could be attributed to increased expression of GLUT4 in rats treated with curcumin. Transformation on cell surface by GLUT4 relies on insulin signaling pathway [29] and IR decreases GLUT4 expression [30]. Not only GLUT4 but also $ER\alpha$ are involved in glucose metabolism. Increased $ER\alpha$ expression promotes the sensitivity of skeletal muscle cells to insulin and causes to consume the glucose [31]. It is clear that curcumin improves IR through involvement in the expression of GLUT4 and $ER\alpha$. Another study showed that curcumin increases the gene expression of GLUT4 but synergistic effects of curcumin and insulin is stronger in comparison to insulin that could be attributed to the competitive activity of insulin and

curcumin in activation of gene expression [32]. It could be speculated that curcumin could improve insulin resistance by influencing on GLUT4 and ER α gene expression. Rats with PCOS showed a number of cystic follicles and decreased corpus luteum. Similar results were reported by previous studies [25]. Oral administration of curcumin could decrease cystic follicles and increase corpus luteum. Increased corpora lutea implicates ovulation and normal estrous cyclicity. Follicles in the different steps of development with oocytes and clear, visible granulosa cell layer were seen in curcumin groups.

Conclusion

In conclusion, our findings show that curcumin has a significant protective effect in patients with PCOS and in addition, the results also show that oral administration of curcumin prevents diabetes. A novel finding of curcumin usage was a significant decrease in insulin resistance and increased expression of GLUT4 and ER α expression that was observed in gene studies and increased folliculogenesis in the ovaries. We can safely advise to use of curcumin for treatment of patients with PCOS and/or as adjuvant treatment.

Ethical Considerations

Compliance with ethical guidelines

Approval for this study was obtained from International Center for Intelligent Research.

Funding

This study was supported by a grant from International Center for Intelligent Research (ICIR-2018-184157).

Authors' contributions

All authors contributed toward data analysis, drafting and revising the paper and agreed to be responsible for all the aspects of this work.

Conflict of interest

The authors declared no conflict of interest.

References

1. Arikawe AP, Iranloye BO, Ogunsola AO, Daramola AO. Chronic fructose consumption as a model of polycystic ovary syndrome in pregnant female sprague-dawley rats. *Afr J Biomed Res.* 2013; 15(1): 7–13.
2. Zhai HL, Wu H, Xu H, Weng P, Xia FZ, Chen Y, Lu YL. Trace glucose and lipid metabolism in high androgen and high-fat diet induced polycystic ovary syndrome rats. *Reprod Biol Endocrinol.* 2012; 10: 5. [\[crossref\]](#)
3. Ghasemzadeh A, Farzadi L, Khaki A, Ahmadi SK. Effect of Allium cepa seeds ethanolic extract on experimental polycystic ovary syndrome (PCOS) apoptosis induced by estradiol-valerate. *Life Sci.* 2013; 10: 170–175.
4. Demirel MA, Ilhan M, Sutar I, Keles H, Akkol EK. Activity of Corylus avellana seed oil in letrozole-induced polycystic ovary syndrome model in rats. *Rev Bras de Farmacogn.* 2016; 26(1): 83–88. [\[crossref\]](#)
5. Makker A, Goel MM, Das V, Agarwal A. PI3K-Akt-mTOR and MAPK signaling pathways in polycystic ovarian syndrome, uterine leiomyomas and endometriosis. An update. *Gynecol Endocrinol.* 2012; 28(3): 175–181. [\[crossref\]](#)
6. Zhao Y, Zhang C, Huang Y, Yu Y, Li R, Li M, Liu N, Liu P, Qiao J. Up-regulated expression of WNT5a increases inflammation and oxidative stress via PI3K/AKT/NF- κ B signaling in the granulosa cells of PCOS patients. *J*

- Clin Endocrinol Metab.* 2015; 100(1): 201–211. [\[crossref\]](#)
7. Skrgatić L, Baldani DP, Gersak K, Cerne JZ, Ferik P, Corić M. Genetic polymorphisms of INS, INSR and IRS-1 genes are not associated with polycystic ovary syndrome in Croatian women. *Coll Antropol.* 2013; 37(1): 141–146. [\[PMID\]](#)
 8. Strowitzki T, von Eye HC, Kellerer M, Haring HU. Tyrosine kinase activity of insulin-like growth factor I and insulin receptors in human endometrium during the menstrual cycle: cyclic variation of insulin receptor expression. *Fertil. Steril.* 1993; 59(2): 315–322. [\[crossref\]](#)
 9. Sheets EE, Tsibris JC, Cook NI, Virgin SD, DeMay RM, Spellacy WN. In vitro binding of insulin and epidermal growth factor to human endometrium and endocervix. *Am J Obstet Gynecol.* 1985; 153(1): 60–65. [\[crossref\]](#)
 10. Mioni R, Chiarelli S, Xamin N, Zuliani L, Granzotto M, Mozzanega B, Maffei P, Martini C, Blandamura S, Sicolo N. Evidence for the presence of glucose transporter 4 in the endometrium and its regulation in polycystic ovary syndrome patients. *J Clin Endocrinol Metab.* 2004; 89(8): 4089–4096. [\[crossref\]](#)
 11. Zhai J, Liu C-x, Tian Z-r, Jiang Q-h, Sun Y-p. Effects of metformin on the expression of GLUT4 in endometrium of obese women with polycystic ovary syndrome. *Biol Reprod.* 2012; 87(2): 307–314. [\[crossref\]](#)
 12. Carvajal R, Rosas C, Kohan K, Gabler F, Vantman D, Romero C, Vega M. Metformin augments the levels of molecules that regulate the expression of the insulin-dependent glucose transporter GLUT4 in the endometria of hyperinsulinemic PCOS patients. *Hum Reprod.* 2013; 28(8): 2235–2244. [\[crossref\]](#)
 13. Hulchiy M, Nybacka A, Sahlin L, Hirschberg AL. Endometrial expression of estrogen receptors and the androgen receptor in women with polycystic ovary syndrome: a lifestyle intervention study. *J Clin Endocrinol Metab.* 2016; 101(2): 561–571. [\[crossref\]](#)
 14. Meyer MR, Clegg DJ, Prossnitz ER, Barton M. Obesity, insulin resistance and diabetes: sex differences and role of oestrogen receptors. *Acta Physiol (Oxf).* 2011; 203(1): 259–69. [\[crossref\]](#)
 15. Quaynor SD, Stradtman EW, Kim HG, Shen Y, Chorich LP, Schreihofner DA, Layman LC. Delayed puberty and estrogen resistance in a woman with estrogen receptor alpha variant. *N Engl J Med.* 2013; 369(2): 164–171. [\[crossref\]](#)
 16. Badawy Ahmed, Elnashar Abubaker. Treatment options for polycystic ovary syndrome. *Int J Womens Health.* 2011; 3: 25–35. [\[crossref\]](#)
 17. Choi SH, Shapiro H, Robinson GE, Irvine J, Neuman J, Rosen B, et al. Psychological side-effects of clomiphene citrate and human menopausal gonadotrophin. *J Psychosom Obstet Gynaecol.* 2005; 26: 93–100. [\[PMID\]](#)
 18. Pari L, Murugan P. Effect of tetrahydrocurcumin on blood glucose, plasma insulin and hepatic key enzymes in streptozotocin induced diabetic rats. *J Basic Clin Physiol Pharmacol.* 2005; 16(4): 257–274. [\[PMID\]](#)
 19. Pari L, Murugan P. Antihyperlipidemic effect of curcumin and tetrahydrocurcumin in experimental type 2 diabetic rats. *Ren Fail.* 2007; 29(7): 881–889. [\[crossref\]](#)
 20. Aggarwal BB, Banerjee S, Bharadwaj U, Sung B, Shishodia S, Sethi G. Curcumin induces the degradation of cyclin E expression through ubiquitin-dependent pathway and up-regulates cyclin dependent kinase inhibitors p21 and p27 in multiple human tumor cell lines. *Biochem Pharmacol.* 2007; 73(7): 1024–1032. [\[crossref\]](#)
 21. Watson JL, Greenshields A, Hill R, Hilchie A, Lee PW, Giacomantonio CA, et al. Curcumin-induced apoptosis in ovarian carcinoma cells is p53-independent and involves p38 mitogen activated protein kinase activation and down regulation of Bcl-2 and survivin expression and Akt signalling. *Mol Carcinog.* 2010; 49(1): 13–24. [\[crossref\]](#)

22. Kadasi A, V.Sirotkin A, Maruniakova N, Kolesarova A, Bulla J, Grossmann R. The effect of Curcumin on secretory activity, proliferation and apoptosis of the porcine ovarian granulose cells. *JMBFS*. 2012; 2(1): 349–357.
23. Kavitha A, Narendra B, Sathish K, Veena K. Evaluation of effects of Commiphora wightii in dehydroepiandrosterone (DHEA) induced polycystic ovary syndrome (PCOS) in rats. *Pharma Tutor*. 2016; 4(1): 47–55.
24. Miladpour B, Rasti M, Owji AA, Mostafavipour Z, Khoshdel Z, Noorafshan A, Zal F. Quercetin potentiates trans differentiation of bone marrow mesenchymal stem cells into the beta cells in vitro. *J Endocrinol Invest*. 2017; 40(5): 513–521. [\[crossref\]](#)
25. Neisy A, Zal F, Seghatoleslam A, Alae S. Amelioration by quercetin of insulin resistance and uterine GLUT4 and ERα gene expression in rats with polycystic ovary syndrome (PCOS). *Reprod Fertil Dev*. 2019; 31(2): 315–323. [\[crossref\]](#)
26. Mesbahzadeh B, Garmsiri M, Jalalvand F, Shojaie L, Kakar MA. Oral administration of menthol could alleviate adverse effects of polycystic ovarian syndrome in blood biochemical parameters and antioxidant status in experimental Wister rats. *GMJ Med*. 2017; 1(1):9–14. [\[crossref\]](#)
27. Diamanti-Kandarakis E, Argyrakopoulou G, Economou F, Kandaraki E, Koutsilieris M. Defects in insulin signaling pathways in ovarian steroidogenesis and other tissues in polycystic ovary syndrome (PCOS). *J Steroid Biochem Mol Biol*. 2008; 109(3–5): 242–246. [\[crossref\]](#)
28. Winkler G, Kiss S, Keszthelyi L, Sa'pi Z, Ory I, Salamon F, Kova'cs M, Vargha P, Szekeres O, Speer G. Expression of tumor necrosis factor (TNF)-α protein in the subcutaneous and visceral adipose tissue in correlation with adipocyte cell volume, serum TNF α, soluble serum TNF-receptor-2 concentrations and C-peptide level. *Eur J Endocrinol*. 2003; 149(2): 129–135. [\[crossref\]](#)
29. Patel N, Huang C, Klip A. Cellular location of insulin triggered signals and implications for glucose uptake. *Pflugers Arch*. 2006; 451(4): 499–510. [\[crossref\]](#)
30. Fornes R, Ormazabal P, Rosas C, Gabler F, Vantman D, Romero C, Vega M. Changes in the expression of insulin signaling pathway molecules in endometria from polycystic ovary syndrome women with or without hyperinsulinemia. *Mol Med*. 2010; 16(3–4): 129. [\[crossref\]](#)
31. Riant E, Waget A, Cogo H, Arnal JF, Burcelin R, Gourdy P. Estrogens protect against high-fat diet-induced insulin resistance and glucose intolerance in mice. *Endocrinol*. 2009; 150(5): 2109–2117. [\[crossref\]](#)
32. Moradi A, Bahrami M, Eslami G, Mohiti-Ardekani J. The Effect of Curcumin on GLUT4 gene expression as a diabetic resistance marker in C2C12 myoblast cells. *Iran J Diabetes Obesity*. 6(2): 98–105.

GMJ Medical Press, LLC

Copyright. © 2018 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation. Shahmoradi M, Askaripour M, Rajabi S, Dzigandzli G. Beneficial Effects of Curcuminon on Rats with Polycystic Ovary Syndrome: Evaluation of the Gene Expression of GLUT4, Er α and Insulin Resistance. GMJ Medicine. 2018; 2: 80–87.

<https://doi.org/10.29088/GMJM.2018.80>